

On-farm Demonstration of Improved Faba Bean Varieties in Bale and West Arsi Zones, Southeastern Oromiya, Ethiopia

Bayeta Gadissa, Ayalew Sida and Amare Biftu

Oromiya Agricultural Research Institute (OARI), Sinana Agricultural Research Center (SARC) P.O.Box-208, Bale-Robe, Ethiopia

Abstract: On-farm demonstration of improved faba bean varieties was conducted in Adaba and Dodola districts of West Arsi Zone and Sinana, Goba and Agarfa of Bale Zone. The main objective of the study was to demonstrate and evaluate recently released (Aloshe) variety along with standard check. The demonstration was under taken on single plot of 10m x 10m area for each variety with the spacing of 40cm between rows and recommended seed rate of 180kg/ha and fertilizer rate of 100qt/ha NPS. Mini-field day involving different stakeholders was organized at each respective site. Yield data per plot was recorded and analysed using descriptive statistics, while farmers' preference to the demonstrated varieties was identified using focused group discussion and summarized using pair wise and simple ranking methods. The demonstration result revealed that Aloshe variety performed better than the standard check (Mosisa variety) with an average yield of 35.97qt/ha, while that of the standard check was 28.5qt/ha. Aloshe variety had 26.21% yield advantage over the standard check. This variety was selected by farmers in Adaba, Dodola and Agarfa. Thus, Aloshe variety was recommended for further scaling up in the area it was selected.

Keywords: Demonstration, Farmers' preference, faba bean, Aloshe, Selection criteria.

INTRODUCTION

Faba bean (*Vicia faba* L) is mainly grown in the highlands (1800-3000 m.a.s.l) of Ethiopia (Yohannes, 2000). Faba bean production ranks the 1st among pulse crops in area and volume of production in the country. From 1,598,806.51 hectares of land allocated for pulse in 2017/2018 production season, faba bean covered 437,106.04 hectares of land from which 9,217,615.35 quintals of grain was produced with the productivity of 21.09 qt/ha (CSA, 2018).

In Bale, 15,347.32 ha of land was covered by faba bean and 372,559.31 quintals of grain was produced with the productivity of 24.28 qt/ha (CSA, 2017). Faba bean has ecological and economic importance and used for food (rich in protein), income source and foreign currency (attractive market price), soil fertility restoration (NP) and food security. Bale and West Arsi Zones are characterized by integrated (mixed) farming systems in which most of the crop areas were under cereal based mono-cropping (Bekele, 2011). Crop diversification can be a means to stay in sustainable crop production in the study zones. Faba bean are the best break crops for wheat production. Bread wheat grown after these crops gave higher grain yield than after cereal crops with a yield advantage of 15% (Sinana ARC Profile, 2014).

Developing high yielding, disease tolerant and stable varieties that can meet increasing food demand of the growing human population, improve the income and livelihood of farmers are

very important. Consequently, **Aloshe variety** has recently released by SARC with yield potential of 35-50qt/ha. The yield advantage of Aloshe over standard and local checks is 20.39 % and 35.08%, respectively.

The two way feedback between farmers and researchers is indeed vital component of high yielder and disease and pest resistant varietal development process (Getachew. *et al.*, 2008). Thus, undertaking participatory demonstration, evaluation, validation and dissemination of improved faba bean varieties with the participation of farmers and other stakeholders for sustainable production and productivity is paramount.

OBJECTIVES

1. To evaluate the yield performance of faba bean varieties under farmers' condition in Bale and West Arsi zones;
2. To create awareness on the importance of faba bean varieties among farmers, DAs, SMSs and other participant stakeholders;
3. To collect farmers' feedbacks on faba bean varieties for further development of faba bean technologies;

METHODOLOGY

Description of the Study Area

The research was carried out at Adaba and Dodola districts of West Arsi Zone and Goba, Sinana and Agarfa districts of Bale zone, Oromiya National Regional State (ONRS), Ethiopia. Bale and west Arsi zones are among the 20 Administrative zones

located in south eastern parts of Oromiya, Ethiopia.

Site and Farmers Selection

On-farm demonstration of improved faba bean varieties was carried out at Adaba and Dodola districts of West Arsi Zone and Goba, Sinana and Agarfa districts of Bale zone. Purposive sampling methods were employed to select the districts based on the potential of the crop. One PA from Adaba, two PAs from Dodola, Goba, Agarfa and Sinana were selected based on accessibility or vicinity to the road. Similarly, two trial farmer from each PA of Adaba and Dodola districts and one trial farmer from each PA of Goba, Sinana and Agarfa districts were used to carry out the demonstration process considering each farmer's field as replication of the trial. The activity was done on a total of 12 trial farmers.

MATERIALS USED AND FIELD DESIGN

Improved faba bean variety (Aloshe) was demonstrated and compared with standard check Mosisa. The demonstration was under taken on simple plot design of 10m x 10m area for each variety. Full packages were applied. In addition, twice hand weeding was done on time.

SARC was the source of all agricultural inputs. Farm preparations were carried out by trial/hosting farmers, whereas land leveling, planting, first and second weeding, follow up and visit, harvesting, threshing were handled and managed by SARC.

Data Collection

Both qualitative and quantitative data were collected using appropriate data collection methods such as direct field observation/measurements, key informant interview and focused group discussion (FGD). Yield data per plot in all locations were recorded. Farmers' preference to the demonstrated varieties was identified.

Data Analysis

Descriptive statistics was used to analyze the yield data. Pair wise ranking and simple matrix ranking were used to compare traits of demonstrated varieties.

RESULT AND DISCUSSION

Training

Training was given to farmers, DAs, and agricultural experts on faba bean crop production techniques and management packages, agro-chemical applications and safety precautions. Stakeholders such as zone and district level agriculture development office, Zone and district level cooperative promotion offices, zone and district level agricultural inputs regulations and quarantine experts were invited and participated during consultation meeting and training.

Yield Performance of Demonstrated Varieties

The mean yield obtained from Adaba, Dodola, Goba, Sinana and Agarfa were summarized in the chart below.

Table 1: yield Performance of demonstrated varieties

No	Variety	Yield obtained (Qt/ha)						Yield advantage over standard check
		Dodola	Adaba	Goba	Sinana	Agarfa	Mean	
1	Aloshe	35.5	34	40.68	36.85	32.8	35.97	26.21%
2	Mosisa	30.5	29.5	27.75	26.14	28.6	28.5	-

The newly released faba bean variety Aloshe performed better than the standard check Mosisa all over the locations. It is the high yielder at all demonstration sites. Aloshe variety gave a mean yield of 35.5qtha⁻¹, 34qt/ha, 40.68qt/ha, 36.85qt/ha and 32.8qt/ha at Dodola, Adaba, Goba, Sinana and Agarfa districts respectively with overall mean yield of 35.97qt/ha. Similarly, Mosisa variety gave

a mean yield of 30.5qt/ha, 29.5qt/ha, 27.75qt/ha, 26.14qt/ha and 28.6qt/ha at Dodola, Adaba, Goba, Sinana and Agarfa districts respectively with overall mean yield of 28.5qt/ha. It also has 26.21% yield advantage over the standard check. The cost benefit ratio reveals that the new variety has higher net profit (1.94) than the standard check (1.38).

Table 2: Cost-Benefit Analysis of the Demonstrated varieties

No	Variables	Varieties	
		Aloshe	Mosisa
	Yield obtained (qt/ha)	35.97	28.50
	Sale price (ETB/qt)	1800	1800
	Gross Returns (Price X Qt) TR	64746	51300
	Total Variable Costs TVC (ETB/ha)	14030	13540
	Fixed cost (FC)	8000	8000
	Total cost (TC)	22030	21540
	Net Return (GR-TC)	42716	29760
	Benefit cost ratio (NR/TVC)	1.94	1.38

Table 3: Result of Independent sample t test

	Test for Equality of Variances		t-test for Equality of Means				
	F	Sig.	T	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference
Equal variances assumed	1.285	.290	4.808	8	.001	7.47	1.55

As shown above (table 3), there was statistically significant difference between the yield of Aloshe and Mosisa varieties. There was also a mean difference of 7.47qt/ha between both varieties.

Farmers' preference of variety traits were identified and presented by pair wise ranking. Accordingly, yield, early maturity, drought tolerance, adaptability to the environment and disease tolerance were the top five priority concern given by farmers (Table 4).

Table 4: Pair wise ranking result to rank variety traits in order of importance

No	variety traits	A	B	C	D	E	F	G	H	I	J	K	Frequency	Rank
1	A												6	5 th
2	B	A											4	6 th
3	C	A	B										4	6 th
4	D	A	D	C									4	6 th
5	E	E	E	E	E								7	4 th
6	F	A	B	C	D	E							2	9 th
7	G	A	B	C	D	E	F						0	11 th
8	H	A	B	C	D	E	F	H					1	10 th
9	I	I	I	I	I	I	I	I	I				8	3 rd
10	J	J	J	J	J	J	J	J	J	J			9	2 nd
11	K	K	K	K	K	K	K	K	K	K	K		10	1 st

A= Disease tolerance, B= Pod/plant, C= Seed/plant, D= Tiller, E= Adaptability, F= Seed size, G= Plant height, H= Marketability, I= Drought tolerance, J= Early maturity, K= Yield

Table 5: Rank of the varieties based on farmers' selection criteria

No	Varieties	Rank	Reasons
1	Aloshe	1 st	High yielder, pod/plant(30), seed/plant(59), more tolerant to disease, tiller(3-4), more adaptable to the environment, more adaptable to the soil, big seed size, good plant height, more marketable, more resistant to drought
2	Mosisa	2 nd	Low yielder, pod/plant(20), seed/plant(46), less tolerant to disease, tiller(3-4), less adaptable to the environment, less adaptable to the soil, small seed size, shorter plant height, less marketable, less resistant to drought

Table 6: Rank of the varieties based on farmers' selection criteria at Sinana district

No	Varieties	Rank	Reasons
1	Aloshe	2 nd	Late mature, tiller (3-4), plant height (long), moderate tolerant to frost, seed/pod, pod/plant, tiller (3-5), but more tolerant to disease
2	Mosisa		Early mature, good plant height (medium height), seed/pod (2-4), pod/plant (15-28),

		1 st	tiller (3-6)
--	--	-----------------	--------------

CONCLUSIONS AND RECOMMENDATIONS

On-farm demonstration and evaluation of faba bean varieties was carried out on twelve (12) representative trial farmers' fields. Improved variety viz. *Aloshe* was demonstrated along with *Mosisa* variety which is the standard check. Accordingly, *Aloshe* gave higher yield than *Mosisa* variety.

Moreover, *Aloshe* was selected by participant farmers in Adaba, Dodola, Goba and Agarfa due to high yielder, pod/plant(30), seed/plant(59), more tolerant to disease, tiller(3-4), more adaptable to the environment, more adaptable to the soil, big seed size, good plant height, more marketable, more resistant to drought. Similarly, farmers selected *Mosisa* variety in Sinana district due to early mature, good plant height (medium height), seed/pod (2-4), pod/plant (15-28), tiller (3-6). Based on these facts, *Aloshe* variety was recommended for further scaling up in the area it was selected.

REFERENCES

1. Bekele, D. "Analysis of Rural Women Farmers' Drudgery and Their Role in Agricultural Production: The Case of Sinana Districts, Bale Zone, Oromiya National Regional State, Ethiopia." *M. Sc. Thesis. Haramaya University, Ethiopia* (2011).
2. Central Statistical Agency (CSA). "The Federal Democratic Republic of Ethiopia. Central Statistical Agency Agricultural Sample Survey 2016/2017 (2009 E.C.): Report on Area and Production of Major Crops (Private Peasant Holdings, Meher Season)." 1 *Addis Ababa, Ethiopia* (2017).
3. Central Statistical Agency (CSA). "The Federal Democratic Republic of Ethiopia. Central Statistical Agency Agricultural Sample Survey 2017/2018 (2010 E.C.): Report on Area and Production of Major Crops (Private Peasant Holdings, Meher Season)." *Addis Ababa, Ethiopia* (2018).
4. Belay, G., Tefera, H., Getachew, A., Assefa, K. and Metaferia, G. "Highly client-oriented breeding with farmer participation in the Ethiopian cereal tef [*Eragrostistef* (Zucc.) Trotter]." *African Journal of Agricultural Research* 3.1 (2018): 022-028.
5. Degago Y. "Faba Bean (*Vicia faba*) in Ethiopia." *Institute of Biodiversity Conservation and Research (IBCR) Addis Ababa, Ethiopia* (2000): 43.

Source of support: Nil; **Conflict of interest:** Nil.

Cite this article as:

Gadissa, B., Sida, A. and Biftu, A. "On-farm Demonstration of Improved Faba Bean Varieties in Bale and West Arsi Zones, Southeastern Oromiya, Ethiopia." *Sarcouncil Journal of Multidisciplinary* 2.3 (2022): pp 4-7